

UNDETECTED SOURCES ALLOW TRANSMISSION OF THE LY α LINE FROM GALAXIES PRIOR TO REIONIZATION

J. STUART B. WYITHE¹ AND ABRAHAM LOEB²
 swyithe@isis.ph.unimelb.edu.au; aloeb@cfa.harvard.edu
Draft version February 2, 2008

ABSTRACT

The discovery of Ly α emission from galaxies at redshifts beyond $z \sim 6.5$ should not be naively interpreted as implying that the intergalactic medium (IGM) had been reionized at higher redshifts. We show that a cluster of faint undetected sources around each observed galaxy generates an HII region sufficiently large to allow transmission of the galaxy's Ly α line prior to reionization. We also show that quasars may contribute a significant fraction of the ionizing photons to HII regions around galaxies with a velocity dispersion larger than $\sim 100 \text{ km s}^{-1}$. These contributing quasars are not usually seen due to the small fraction of time they spend in a luminous phase.

Subject headings: cosmology: theory - galaxies: formation

1. INTRODUCTION

The identification of a Gunn-Peterson trough in the spectra of the most distant quasars at redshifts of $z \sim 6.3$ – 6.4 (Fan et al. 2004) hints that the reionization of cosmic hydrogen was completed only around that time, about a billion years after the big bang (White et al. 2003; Wyithe & Loeb 2004). On the other hand, the recent discovery of Ly α emission from galaxies beyond a redshift of $z \sim 6.5$ (Hu et al., 2002; Kodaira et al., 2003; Rhoads et al. 2004; Kurk et al. 2004; Stern et al. 2004) was naively interpreted as an indication that the surrounding intergalactic medium (IGM) was highly ionized, in apparent conflict with the quasar data. Although a galaxy is often surrounded by an HII region of ionized hydrogen (Haiman 2002; Santos 2004), its Ly α line could be heavily suppressed by the red damping wing of the resonant Ly α absorption in the surrounding IGM (Loeb, Barkana & Hernquist 2004; Cen, Haiman & Messinger 2004; Ricotti et al. 2004). Clustering of sources may result in HII regions that are larger than expected for a single galaxy, rendering the optical depth from the red damping wing of the IGM irrelevant (Gnedin & Prada 2004; Furlanetto, Hernquist & Zaldarriaga 2004). Here we show that undetected sources such as short-lived quasars or a cluster of faint galaxies, are expected to dominate the growth of a galaxy's HII region and allow transmission of its Ly α photons prior to reionization.

2. THE SIZE OF IONIZED REGIONS

The average mean-free-path for an ionizing photon in a fully neutral IGM is much smaller than the size of a cosmological HII region. By balancing the number of ionizing photons and neutral atoms we can therefore estimate the characteristic physical size of a spherical HII region surrounding a galaxy with a dark matter halo of mass M_{halo} (Loeb et al. 2004)

$$D_{\star} = 0.75 \left(\frac{x_{\text{HI}}}{0.1} \right)^{-1/3} \left(\frac{M_{\text{halo}}}{10^{10} M_{\odot}} \right)^{1/3} \left(\frac{f_{\star} f_{\text{esc}} / N_{\text{reion}}}{0.003} \right)^{1/3}$$

¹ University of Melbourne, Parkville, Victoria, Australia

² Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

$$\times \left(\frac{N_{\star}}{4000} \right)^{1/3} \left(\frac{1+z}{7.5} \right)^{-1} \text{ Mpc}, \quad (1)$$

where $f_{\star}$ is the star formation efficiency and f_{esc} is the fraction of ionizing photons that escape from the galaxy into the surrounding IGM of neutral hydrogen fraction x_{HI} . Here we have assumed that the number of ionizing photons per baryon incorporated into (Pop-II) stars is $N_{\star} \sim 4000$ (Bromm, Kudritzki & Loeb 2001), and that N_{reion} photons are required per baryon for the reionization of the HII region. The value of N_{reion} depends on the number of recombinations and hence on the clumpiness of the IGM. The halo mass M_{halo} is a function of its velocity dispersion σ [see Eqs. 22–25 in Barkana & Loeb (2001)], which we set equal to $v_c / \sqrt{2}$, where v_c is the halo circular velocity.

Similarly, the physical size of the HII region generated by an accreting supermassive black-hole of mass M_{bh} is (White et al., 2003)

$$D_{\text{q}} = 0.43 \left(\frac{x_{\text{HI}}}{0.1} \right)^{-1/3} \left(\frac{M_{\text{bh}}}{10^5 M_{\odot}} \right)^{1/3} \left(\frac{\dot{N}_{\gamma,5} / N_{\text{reion}}}{3 \times 10^{53} \text{ s}^{-1}} \right)^{1/3} \times \left(\frac{f_{\text{dyn}}}{0.035} \right)^{1/3} \left(\frac{1+z}{7.5} \right)^{-1} \text{ Mpc}, \quad (2)$$

where $\dot{N}_{\gamma,5}$ is the production rate of ionizing photons during the luminous quasar phase for a black hole of mass $M_{\text{bh}} = 10^5 M_{\odot}$, and f_{dyn} is the fraction of the halo dynamical time during which the quasar shines at its peak luminosity.

There is ample observational data on the number counts and clustering of quasars. We therefore compute the size of quasar HII regions using a simple, physically-motivated, and highly successful model for high redshift quasars that has been developed in earlier papers (Wyithe & Loeb 2003a, 2004). The slow evolution of the quasar clustering correlation length with redshift (Croom et al. 2001) implies that the black-hole mass scales with halo velocity dispersion independent of redshift. This conclusion is supported by direct observations of the M_{bh} – σ relation (Shields et

al. 2003). The observed scaling and its redshift dependence [$M_{\text{bh}} \propto \sigma^5 \propto M_{\text{halo}}^{5/3} (1+z)^{3/2}$] follow naturally from a scenario where black-hole growth is limited by feedback over the galaxy dynamical time (Silk & Rees 1998; Wyithe & Loeb 2003a). By associating quasar activity with major mergers, it has been shown (Wyithe & Loeb 2003a) that when the quasar lifetime equals the galaxy dynamical time the observed number counts of quasars are accurately reproduced over a wide range of redshifts, $1.5 \lesssim z \lesssim 6.5$. The size of the quasar generated HII region (D_q) may therefore be evaluated using equation (2), with a black hole mass that is related to the halo velocity dispersion (Wyithe & Loeb 2003a) through $M_{\text{bh}} = 10^5 (\sigma/54 \text{ km s}^{-1})^5 M_\odot$, and a lifetime that is governed by a value of $f_{\text{dyn}} = 0.035$ corresponding to the typical fraction of the virial radius occupied by the luminous component of the galaxy. We point out that while the association of a total quasar luminosity with a dark-matter halo must be made via a theoretical model, the model used here correctly predicts the number density of high redshift quasars over a wide range of luminosities and redshifts. The total number of ionizing photons produced by a quasar (with a universal spectrum) during its lifetime, reflects a fixed fraction of its black-hole rest-mass energy.

The ratio between the number of ionizing photons produced by the central quasar and by stars is

$$\Gamma_{\text{halo}} = 1.4 \left(\frac{\sigma}{100 \text{ km/s}} \right)^2 \left(\frac{f_{\text{dyn}}}{0.035} \right) \left(\frac{f_\star f_{\text{esc}}}{0.003} \right)^{-1} \left(\frac{N_\star}{4000} \right)^{-1} \times \left(\frac{\dot{N}_{\gamma,5}}{3 \times 10^{53} \text{ s}^{-1}} \right) \left(\frac{1+z}{7.5} \right)^{3/2}. \quad (3)$$

Since the dependence of Γ_{halo} on halo mass is steeper than linear for $f_\star = \text{const}$, quasars become increasingly more important for higher mass galaxies. Given that both massive stars and quasars produce ionizing photons in short-lived episodes, the value of Γ_{halo} reflects their net yield of ionizing photons. The HII region generated by quasars remains as a fossil after the quasar activity ends, since the recombination time is longer than the Hubble time at the mean IGM density for $z \lesssim 8$. Hence, relic HII regions from prior quasar activity must be considered when estimating the transmission of Ly α photons from high redshift galaxies. In what follows we show that this remains true even when considering the enhanced recombination rate of the overdense IGM around massive galaxies.

3. STRÖMGREN SPHERES AT HIGH-REDSHIFT

Prior to reionization, galaxies generate their own Strömgren (HII) spheres with characteristic sizes given by equations (1) and (2). These spheres expand into a partially neutral IGM. If the Strömgren sphere is produced by a massive galaxy, then the external neutral fraction reflects the mean volume filling factor of all small HII regions produced by its neighboring low-mass galaxies. The clustering of galaxies around a bright galaxy could lead to overlap of their individual HII regions, even if the rest of the IGM is predominantly neutral. As a result the HII region surrounding an observed galaxy could be larger than expected from equations (1) and (2) due to a cluster of faint galaxies that are below the detection threshold of the observations. The physical size R_s of the HII region

surrounding a galaxy of mass M_{halo} can be determined self-consistently with the mean neutral fraction into which the HII region expands, through the relation

$$R_s^3 = D_{\text{halo}}^3 + \int_{R_{\text{vir}}}^{R_s} 4\pi R^2 dR \int_{M_{\text{min}}}^{M_{\text{halo}}} dM \frac{D_M^3}{\Delta_R^2} \frac{dn(R, M_{\text{halo}})}{dM}, \quad (4)$$

where the sizes of the Strömgren spheres around the central halo of mass M_{halo} and a galaxy of mass M , respectively, are $D_{\text{halo}}^3 = D_{\text{halo,q}}^3 + D_{\text{halo},\star}^3$ and $D_M^3 = D_{M,q}^3 + D_{M,\star}^3$. Equation (4) accounts for the higher IGM density due to infall around a massive halo through the nonlinear density contrast $\Delta_R(R_s, M_{\text{halo}}) = \rho/\bar{\rho}$ (where $\bar{\rho}$ is the mean background density) at a radius R from a halo of mass M_{halo} (Barkana 2004). The associated density profile reproduces the appropriate two-point correlation function in the linear regime (Barkana 2004) and is evolved into the nonlinear regime in spherical shells. The number of galaxies surrounding the central halo is found from the Press-Schechter (1974) mass function of halos, which in the vicinity of a massive halo must be modified relative to the background universe. Within the spherical infall model, shells with density ρ at radius R behave as if they are embedded in a closed universe with a matter density parameter $\Omega_m(R) = \rho/\rho_c$, where $\rho_c = 3(\dot{R}/R)^2/(8\pi G)$ and $\dot{R}$ is the physical radial velocity of the shell (the cosmological constant is negligible at the high redshifts of interest.) Thus, $dn(R, M_{\text{halo}})/dM$ is the physical number density of halos per unit mass computed in a universe with a density parameter $\Omega_m(R)$ (with a normalization and a growth factor based on the local shell density). Each HII region reaches its final comoving volume when the production rate of ionizing photons is balanced by the recombination rate interior to that volume. In this phase, the HII volume is inversely proportional to the square of the mean density times a clumping factor. The clumping is accounted for by N_{reion} , but an additional factor of Δ_R^{-2} needs to be included on the right-hand-side of equation (4) to account for recombinations at the mean density within each shell. We assume that the most massive galaxy interior to the HII region is the one observed. This conservative assumption is reflected in the upper limit of the mass integration. The value M_{min} is the minimum mass (Barkana & Loeb 2001), below which gas infall is suppressed from an ionized region that has been heated to 10^4 K .

Next we specify the free parameters in our model. The absorption spectra of the highest redshift quasars indicate that the most recent reionization of hydrogen occurred at $z \sim 6$ (Fan et al. 2001; Wyithe & Loeb 2004). Moreover, there is evidence for a temperature rise in the Ly α forest associated with the reionization of HeII at $z \sim 3.5$ (Theuns et al. 2002). We set the free parameters by requiring that hydrogen be reionized by $z = 6$, and helium be doubly reionized by $z = 3.5$. Population-II stars do not have spectra that are sufficiently hard to ionize HeII, and so in the absence of Population-III stars at low redshifts the ionization fraction of HeII is determined by quasars. The global ionization fractions of intergalactic hydrogen and helium are therefore

$$F_{\text{ion,HII}} = \int_{M_{\text{min}}}^{\infty} dM \frac{dn}{dM} \frac{4\pi}{3} (D_{M,q}^3 + D_{M,\star}^3) \quad \text{and}$$




FIG. 1.— The physical size of cosmological HII regions, and the detectability of high redshift galaxies. *Left*: The neutral fraction of the intergalactic hydrogen vs. redshift [see equation (5)], with the constraints that hydrogen is reionized at $z = 6$ and that helium is doubly-reionized at $z = 3.5$. The gray, left-most curve shows the neutral fraction of HeII. *Center*: The size of an HII region at ($z = 7$) due to clustered faint sources around a central galaxy with a velocity dispersion σ . Also shown (gray curves) are values for the naive estimate of the size of the HII region, assuming a fully neutral IGM at mean density [Eq. (1)]. *Right*: Optical depth for Ly α absorption by the damping wing of the distant IGM, at the center of an HII region around a galaxy with a velocity dispersion σ . From left to right, the different sets of lines correspond to redshifts $z = 6.5$, $z = 7$ and $z = 10$, respectively. The different line types mark the four cases **A–D** (see main text) labeled in the middle panel.

$$F_{\text{ion,HeIII}} = \int_{M_{\text{min}}}^{\infty} dM \frac{dn}{dM} \frac{4\pi}{3} D_{\text{M,He,q}}^3. \quad (5)$$

where $D_{\text{M,He,q}} \approx 0.85 D_{\text{M,q}}$ assuming the median spectral energy distribution of quasars (Telfer et al. 2002), and that $\sim 50\%$ of the photons of sufficient energy to double-ionize helium are utilized in the reionization of hydrogen (Wyithe & Loeb 2003b). The ratio $\dot{N}_{\gamma,5}/N_{\text{reion}}$ may be determined from the requirement that $F_{\text{ion,HeIII}}$ approach unity at $z \sim 3.5$. Assuming, for simplicity, that $N_{\text{reion}} = \text{const}$ and that the spectral slope of quasars blueward of the ionization threshold of hydrogen is specified by their median spectral template (Telfer et al. 2002), the ratio $f_{\star} f_{\text{esc}}/N_{\text{reion}}$ can then be determined from the requirement that $F_{\text{ion,HII}}$ approach unity at $z \sim 6$. Gas clumping is expected to increase with cosmic time. By calibrating N_{reion} based on the reionization of helium and hydrogen at $z \sim 3.5$ and 6, we therefore overestimate the significance of recombinations at earlier times and derive conservative results for the minimum transmission of the Ly α line. We note that even though our model is calibrated by requiring that hydrogen reionization be completed at $z \sim 6$ by Pop-II stars and quasars, an early generation of massive Pop-III stars may be required to produce a first partial or full reionization of hydrogen at $z \sim 15\text{--}20$. An early reionization would in provide consistency between the quasar absorption at $z \sim 6$ and the large scale polarization anisotropies of the microwave background detected by the *Wilkinson Microwave Anisotropy Probe* (WMAP; Wyithe & Loeb 2003b; Cen 2003).

Reionization histories may now be calculated based on equation (5) for different star formation prescriptions, with $\dot{N}_{\gamma,5}$ and $f_{\star}$ adjusted to meet the above global requirements. We adopt feedback-regulated prescriptions (Dekel & Woo 2002; Wyithe & Loeb 2003a) in which the star formation efficiency in a galaxy depends on its velocity dispersion σ , with $f_{\star} = 0$ for $\sigma < \sigma_{\text{min}}$, $f_{\star} = f_{\star,\text{crit}} (\sigma/\sigma_{\text{crit}})^2$ for $\sigma_{\text{min}} < \sigma < \sigma_{\text{crit}}$, and $f_{\star} = f_{\star,\text{crit}}$ for $\sigma > \sigma_{\text{crit}}$. We consider four cases in this paper: **(A)** $\sigma_{\text{min}} = \sigma_{\text{crit}} = 10 \text{ km s}^{-1}$; **(B)** $\sigma_{\text{min}} = 10 \text{ km s}^{-1}$, $\sigma_{\text{crit}} = 120 \text{ km s}^{-1}$; **(C)** $\sigma_{\text{min}} = \sigma_{\text{crit}} = 50 \text{ km s}^{-1}$; and **(D)** $\sigma_{\text{min}} = 50 \text{ km s}^{-1}$, $\sigma_{\text{crit}} = 120 \text{ km s}^{-1}$. The values of $\sigma_{\text{min}} = 50 \text{ km s}^{-1}$, and $\sigma_{\text{min}} = 10 \text{ km s}^{-1}$, correspond to the threshold for infall of gas from a photo-ionized IGM and the threshold

for atomic hydrogen cooling, respectively. The value of $\sigma_{\text{crit}} = 120 \text{ km s}^{-1}$ is calibrated empirically based on the local sample of galaxies (Kauffmann et al. 2003; Dekel & Woo 2002). We also examine the case of $\sigma_{\text{crit}} = \sigma_{\text{min}}$ to reflect our ignorance about whether a similar feedback operates at high redshifts, when the dynamical time within galaxies becomes comparable to the lifetime of massive stars. We have adopted the set of cosmological parameters determined by WMAP (Spergel et al. 2003), and find that a constraint of late reionization at $z \sim 6$ requires values for the parameter combination $N_{\star} f_{\star,\text{crit}} f_{\text{esc}}/N_{\text{reion}}$ of 11, 2, 22, and 17 for models **A–D**.

In the central panel of Figure 1, we plot the Strömgren radius R_s as a function of the velocity dispersion σ of a central galaxy at a redshift of $z = 7$. The four curves correspond to cases **A–D**. These calculations of R_s are self-consistent with the average neutral fraction of the IGM, which is obtained from the reionization histories (Eq. 5), and plotted in the left panel of Figure 1. A rapid, stellar reionization that completes by $z = 6$ would imply a neutral fraction of $x_{\text{HI}} \sim 0.2\text{--}0.5$ at $z \sim 6.5\text{--}7$, consistent with the analysis of the HII regions around the highest redshift quasars (Wyithe & Loeb 2004). At $z = 7$, the HII regions approach a radius of $R_s \sim 0.5\text{--}1 \text{ Mpc}$ for $\sigma \sim 100 \text{ km s}^{-1}$, comparable to the values inferred for recently discovered galaxies (Kneib et al. 2004; Rhoads et al. 2004). Galaxies with $\sigma \sim 300 \text{ km s}^{-1}$, corresponding to the hosts of the luminous $z \gtrsim 6$ quasars, have $R_s \sim 3\text{--}4 \text{ Mpc}$ in agreement with the observed sizes of their Strömgren spheres (White et al. 2003). For comparison, we also show by gray lines the naive estimate for the size of the HII region using only the stars within the central halo, and assuming a fully neutral IGM at the mean density (Eq. (1)).

4. SUPPRESSION OF THE LY α EMISSION LINE

At $z \sim 6.5\text{--}7$, the ionized regions around galaxies with $\sigma \gtrsim 100 \text{ km s}^{-1}$ are sufficiently large to allow transmission of their Ly α emission line despite the red damping wing of Ly α absorption from the surrounding IGM. It is therefore not surprising that these galaxies are observed, even if they reside in an IGM that is significantly neutral globally. The optical depth τ_{damp} to Ly α absorption at the galaxy redshift (Miralda-Escude 1998) is plotted as a




FIG. 2.— Comparisons between the contributions to the HII region of stars relative to quasars, and of the central galaxy relative to the surrounding cluster of galaxies. *Left*: The ratio between the number of ionizing photons contributed to the HII region by quasars and by stars as a function of the velocity dispersion of the host galaxy [Eq. (7)]. *Right*: The fraction of the HII region volume contributed by the central galaxy as a function of the velocity dispersion of the host galaxy. In each panel the different line types correspond to cases **A–D** as in Figure 1 for galaxies at $z = 7$.

function of σ in the right hand panel of Figure 1 for galaxies at $z \sim 6.5$, $z \sim 7$ and $z \sim 10$. Here the size of the HII region is based on equation (4), while the neutral fraction outside the HII region was computed from equation (5). At $z \sim 10$, a galaxy must possess $\sigma \gtrsim 200 - 300 \text{ km s}^{-1}$ in order to have $\tau_{\text{damp}} < 1$. However at $z = 6.5 - 7$, $\tau_{\text{damp}} \lesssim 1$ for galaxies with $\sigma \gtrsim 70 - 100 \text{ km s}^{-1}$. Hence, observations of galaxies at $z \sim 6.5 - 7$ (Kodaira et al., 2003; Rhoads et al. 2004; Kurk et al. 2004; Stern et al. 2004) are fully consistent with reionization completing at $z \sim 6$. The requirement that HII and HeIII be reionized at $z \sim 6$ and $z \sim 3.5$ respectively, make this conclusion robust against variations in $f_{\star, \text{crit}}$ or N_{reion} .

5. CONTRIBUTIONS TO THE IONIZING FLUX

We now explicitly examine the individual contributions from quasars and from the clustering of faint sources to the HII region around massive galaxies. First, we find the ratio between the quasar and stellar fluxes within the HII region, Γ_{tot} . The flux ratio Γ_{halo} from the central halo is obtained from equation (3). The mean flux ratio $\langle \Gamma_{\text{gal}} \rangle$ from other galaxies within the HII region of radius R_s is then

$$\langle \Gamma_{\text{gal}} \rangle = (R_s^3 - D_{\text{halo}}^3)^{-1} \times \int_{R_{\text{vir}}}^{R_s} 4\pi R^2 dR \int_{M_{\text{min}}}^{M_{\text{halo}}} dM \frac{D_M^3}{\Delta_R^2} \frac{dn(M_{\text{halo}}, R)}{dM} \Gamma, \quad (6)$$

and the average flux ratio within the HII region is therefore

$$\Gamma_{\text{tot}} = \frac{(R_s^3 - D_{\text{halo}}^3) \langle \Gamma_{\text{gal}} \rangle + D_{\text{halo}}^3 \Gamma_{\text{halo}}}{R_s^3}. \quad (7)$$

The left-hand-panel of Figure 2 shows $\Gamma_{\text{tot}}(\sigma)$ for cases **A–D** at $z = 7$. Quasars make a substantial contribution to the Strömgren volume around massive galaxies, particularly in case **A**. However the cluster of smaller galaxies surrounding the central halo contributes an overall ionizing luminosity that is mostly stellar, resulting in a value of Γ_{tot} that is smaller than Γ_{halo} .

The fraction of the HII region excavated by the central galaxy is found from the ratio D_{halo}^3/R_s^3 . This quantity is shown in the right panel of Figure 2 as a function of σ for galaxies at $z = 7$. The regions around massive galaxies are predominantly ionized by the surrounding cluster of smaller galaxies. The luminous quasars discovered at $z \gtrsim 6$ have, during their active phase, luminosities in excess of the time-averaged luminosity of these clusters, and are therefore observed as the source of their HII region.

6. SUMMARY

The massive galaxies discovered prior to reionization should have Strömgren spheres that are much larger than naively expected from the UV continuum flux of their stars. This increase arises for two reasons: (i) there is an additional contribution from earlier quasar activity within the central and surrounding galaxies, and (ii) there is a biased clustering of faint sources surrounding the central galaxy which results in overlap of nearby HII regions at a time before overlap is achieved throughout the universe. The large extent of the HII regions allow transmission of Ly α photons from galaxies. We conclude that the discovery of Ly α emission from bright galaxies at $z > 6.5$ does not indicate a highly ionized IGM. Nevertheless, the strength of the Ly α damping increases as the galaxy luminosity decreases or the redshift increases (see the right-hand panel of Fig. 1). A neutral IGM would therefore suppress the low-luminosity end of the Ly α luminosity function of galaxies at high redshifts.

We thank Daniel Stern for comments on the manuscript. This work was supported in part by NASA grant NAG 5-13292, and by NSF grants AST-0071019, AST-0204514 for AL.

REFERENCES

- Barkana, R., Loeb, A. 2001, Phys. Rep., 349, 125
 Barkana, R. 2004, MNRAS, 347, 57
 Bromm, V., Kudritzki, R., Loeb, A. 2001, ApJ 522, 464
 Cen, R., Haiman, Z., Messinger, A. 2004, ApJ, submitted, astro-ph/403419
 Cen, R. 2003, ApJ, 591, L5
 Croom, S. et al. 2001, MNRAS, 325, 483

- Dekel, A., Woo, J. 2002, MNRAS, 344, 1131
- Fan, X., et al. 2004, Astron. J, in press, astro-ph/0405138
- Furlanetto, S., Hernquist, L. & Zaldarriaga, M. 2004, MNRAS, submitted, astro-ph/0406131
- Gnedin, N.Y., Prada, F., 2004, ApJ, 608, L77
- Haiman, Z. 2002, ApJ, 576, L1
- Hu, E.M., et al. 2002, ApJ, 568, L75
- Kauffman, G., et al. 2003, MNRAS, 341, 54
- Kneib, J.P., et al. 2004, ApJ, in press, astro-ph/0402319
- Kodaira et al. 2003, PASJ, 55, L17
- Kurk, J. D., et al. 2004, A & A Lett., in press, astro-ph/0406140
- Loeb, A., Barkana, R., Hernquist, L. 2004, ApJ., submitted, astro-ph/0403193
- Miralda-Escude, J. 1998, ApJ. 501, 15
- Pello, R., et al. 2004, Astron. Astrophys., 416, L35
- Press, W., Schechter, P. 1974, ApJ, 187, 425
- Rhoads, J.E., et al. 2004 ApJ, in press, astro-ph/0403161
- Ricotti, M., Haehnelt, M.G., Pettini, M., Rees, M.J. 2004, astro-ph/0403327
- Santos, M. 2004, MNRAS, 349, 1137
- Silk, J., Rees, M.J. 1998, A & A, 331, 1
- Shields, G.A., et al. 2003, ApJ 583, 124
- Spergel, D. N., et al. 2003, AJ Supp., 148, 175
- Stern, D., Yost, S. A., Eckart, M. E., Harrison, F. A., Helfand, D., & Djorgovski, S. G. 2004, ApJ, submitted
- Telfer, R. C., Zheng, W., Kriss, G. A., Davidsen, A. F. 2002, AJ, 565, 773
- Theuns, T., Schaye, J., Zaroubi, S., Kim, T.S., Tzanavaris, P., Carswell, B. 2002, ApJ, 567, L103
- White, R., Becker, R., Fan, X., Strauss, M. 2003, AJ, 126, 1
- Wyithe, J.S.B, Loeb, A. 2003a, ApJ, 595, 614
- Wyithe, J.S.B., Loeb, A. 2003b, ApJ 586, 693
- Wyithe, J. S. B., Loeb, A. 2004, Nature, 427, 815
- Wyithe, J.S.B., Loeb, A. 2004, ApJ, submitted, astro-ph/0403614